

Assignment 3  
Intro to Modern Analysis

1. Let  $X$  be a set and consider the discrete metric  $d$  on  $X$  defined by

$$d(p, q) = \begin{cases} 1 & p \neq q \\ 0 & p = q \end{cases}.$$

Show that  $X$  is compact if and only if  $X$  is finite.

2. Let  $x_n$  be the sequence of real numbers

$$x_n = \sqrt{1 + \frac{1}{n}}.$$

(a) Show that  $x_n$  converges to 1.

(b) Calculate

$$\lim_{n \rightarrow \infty} \sqrt{n^2 + n} - n.$$

3. Let  $s_n$  be a sequence of real numbers. Construct a new sequence  $\sigma_n$  by the averages

$$\sigma_n = \frac{s_1 + s_2 + \cdots + s_n}{n}.$$

(a) If  $s_n$  converges to  $s$ , show that  $\sigma_n$  converges to  $s$ .

(b) Construct a sequence  $s_n$  which does not converge but which satisfies  $\sigma_n \rightarrow 0$ .

4. Construct a Cauchy sequence in  $\mathbb{Q}$  that does not converge (to a point of  $\mathbb{Q}$ ). (Hint: It might be useful to use the construction from Example 1.1 of Rudin or Proposition 2 from the notes.)

5. Let  $(X, d)$  be a metric space, and let  $p_n, q_n$  be two Cauchy sequences in  $X$ . Show that the sequence  $d(p_n, q_n)$  is Cauchy in  $\mathbb{R}$ .

6. (Not to be graded!) For a real number  $p \geq 1$ , let  $\ell^p$  denote the vector space of sequences  $x = (x_1, x_2, \dots)$  of real numbers such that

$$\sum_{n=1}^{\infty} |x_n|^p$$

converges.

(a) Show that if  $p \leq q$ , then  $\ell^p \subset \ell^q$ .

(b) Suppose  $p < q$ . Find a sequence  $x \in \ell^q$  but not in  $\ell^p$ .

7. Let  $m$  denote the metric space whose elements are bounded infinite sequences of real numbers together with the metric

$$d(x, y) = \sup_{n=1,2,\dots} |x_n - y_n|.$$

Show that  $m$  is complete.

8. Let  $a_n, b_n$  be sequences of real numbers. Suppose that

- (i)  $\sum_n a_n$  converges
- (ii)  $b_n$  is bounded
- (iii)  $b_n$  is monotonic.

Show that  $\sum_n a_n b_n$  converges.

9. This problem has two parts.

- (a) Show that if  $x, y$  are nonnegative real numbers, then

$$xy \leq \frac{1}{2}(x^2 + y^2).$$

- (b) Let  $a_n$  be a sequence of nonnegative real numbers, and let  $b_n = \sqrt{a_n}/n$ . Show that if  $\sum_n a_n$  converges, then  $\sum_n b_n$  converges.

10. State and prove the convergence or divergence of  $\sum_n a_n$  if

- (a)  $a_n = \sqrt{n+1} - \sqrt{n}$
- (b)  $a_n = \frac{\sqrt{n+1} - \sqrt{n}}{n}$
- (c)  $a_n = (\sqrt[n]{n} - 1)^n$ .

**Extra. (Not to be graded)** Let  $(M, d)$  be a metric space. Let  $\mathcal{M}$  be the set of Cauchy sequences in  $M$ , that is, an element  $p$  of  $\mathcal{M}$  consists of a sequence  $P = (p_1, p_2, p_3, \dots)$  of points of  $M$ .

- (a) Problem 5 shows that we can associate to any two  $P, Q \in \mathcal{M}$  a real number  $\Delta(p, q)$  defined by

$$\Delta(P, Q) = \lim_{n \rightarrow \infty} d(p_n, q_n).$$

Show that the resulting function  $\Delta : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$  is symmetric and nonnegative.

- (b) Define a relation  $\sim$  on  $\mathcal{M}$  by  $P \sim Q$  if and only if  $\Delta(P, Q) = 0$ . Show that the relation  $\sim$  is an equivalence relation on  $\mathcal{M}$ .
- (c) Let  $M^*$  denote the set  $\mathcal{M}/\sim$  of equivalence classes. Show that  $\Delta$  induces a well-defined map on  $M^*$  which is a metric.

- (d) Show that the resulting metric space  $M^*$  is complete.
- (e) For any point  $p \in M$ , let  $P_p$  be the sequence whose terms are all  $p$ . Show that for any two points  $p, q$  in  $M$ , we have

$$\Delta(P_p, P_q) = d(p, q).$$

Conclude that there is a distance-preserving map  $\varphi : M \rightarrow M^*$ .

- (f) Show that  $\varphi(M)$  is dense in  $M^*$ .
- (g) Show that  $\varphi(M) = M^*$  if and only if  $M$  is complete.

As a result of this exercise, we may finally define  $\mathbb{R}$ : we may set  $\mathbb{R} = M^*$  for  $M = \mathbb{Q}$ .